

**Dispatch**

**Wear GLOVES and EYE PROTECTION on ALL CALLS**

**ALL Medical Calls**

Apply **MASK** to crew  
(Surgical mask or N95)

**NON-Medical (Trauma, falls, etc)**

↓

**ASSESS all patients from > 6 feet away**

- Cough or Shortness of Breath
- Sore Throat
- Fever
- Malaise
- "Not feeling well"
- Flu symptoms

**ALL Medical Calls**

**NON-Medical (Trauma, Psych, etc.)**

If **NO**

No patient mask necessary; Crew should maintain their mask

If **YES** to any of the above

Apply **MASKS** to crew, if not already wearing, then approach patient

If **NO**

Use **standard PPE** precautions

**Patient must be SOURCE CONTROLLED** by placing a surgical mask on them.  
Maintain surgical mask prior to arrival at EMS Triage.  
*If unable to do so due to facial features or medical procedures, prenotify receiving ahead of time and expect decon*

**ANY Aerosolizing Procedures**

- Nebulizer treatment
- CPAP
- Intubation
- SGA (King/iGel)
- etc.

Crew must don **GOWNS** and **N95**

**Driving?**

1. **DON** all appropriate PPE on scene
2. **DOFF** appropriately into biohazard bag **BEFORE** entering ambulance cab
3. Have PPE accessible from **exterior** of ambulance
4. **DON** appropriate PPE again upon arrival at the hospital

**After EVERY Call**

1. After **transfer of care**, appropriately **DOFF** all PPE
  - a. Place in biohazard bag
  - b. Dispose of bag appropriately when full
2. Don't **REUSE** PPE between patients
3. **WIPE DOWN** all patient care surfaces after every call
4. **WASH HANDS** thoroughly with soap and water

**DONNING PPE**

1. GOWN 
2. MASK 
3. GOGGLES 
4. GLOVES 

**DOFFING PPE**

1. GLOVES 
2. GOWN 
3. GOGGLES 
4. MASK 