

ASPIRIN

Strong Adult Medical Control: (585) 271-2719

INDICATIONS: ACUTE CORONARY SYNDROME - Chest pain, suspicious of cardiac origin, including signs and symptoms of: retrosternal pain, heaviness, tightness or discomfort in the chest, radiation of pain or discomfort into jaw, arms or shoulders.

CONTRAINDICATIONS: Known Allergy (Hypersensitivity)
Active GI Bleeding

CAUTIONS: GI Bleeding and upset
Already taken prior to EMS arrival

ADVERSE EFFECTS: Heartburn
Nausea / Vomiting
Wheezing

HOW SUPPLIED: 81 mg tablets
(TYPICALLY)

DOSAGES: **ADULT (>30 kg or 66 lbs)** **PEDIATRIC (<30 kg or 66 lbs)**
324 mg (4 x 81 mg tablets) NOT INDICATED

INDICATIONS: If symptoms of shock, airway swelling or respiratory distress are present as the result of an allergic reaction, the provider may:
1. Assist the patient in administering the kit's contents
(the patient has their own anaphylaxis kit)
2. Administer the Auto-Injector
(the agency has completed registration as an EpiPen agency and the provider has been trained in its use)

CONTRAINDICATIONS: None in the presence of anaphylaxis

CAUTIONS: Medical Control **must be contacted** if the patient is not prescribed an Epinephrine Auto-Injector. If Epinephrine has been given, ALS must transport with the patient.
(Do not delay transporting the patient while waiting for ALS to arrive at the scene)

ADVERSE EFFECTS: Tachycardia / Palpitations Dizziness Nausea/Vomiting
Chest Pain Hypertension Nervousness / Anxiety
Headache Tremors

HOW SUPPLIED: Epinephrine Auto-Injector only (Patient prescribed or EMS service authorized)
ADULT: 0.3 mg (EpiPen) PEDIATRIC: 0.15mg (EpiPen Jr)
(TYPICALLY)

DOSAGES: **ADULT (>30 kg or 66 lbs)** **PEDIATRIC (<30 kg or 66 lbs)**
One (1) Adult Auto Injector (0.3 mg Epi) One (1) Pediatric Auto Injector (0.15 mg Epi)

EPINEPHRINE (Auto-Injector)
Strong Adult Medical Control: (585) 271-2719
Strong Pediatric Medical Control: (585) 756-3430

ALBUTEROL (Proventil, Ventolin)

Strong Adult Medical Control: (585) 271-2719
Strong Pediatric Medical Control: (585) 756-3430

INDICATIONS: Signs and symptoms of respiratory distress and bronchospasm/wheezing associated with asthma

CONTRAINDICATIONS: Known Allergy (Hypersensitivity)

CAUTIONS: Patient must have been prescribed a hand held aerosol inhaler
If patient has not previously had albuterol, contact medical control **prior** to administration

ADVERSE EFFECTS:

Tachycardia / Palpitations	Dizziness
Hypertension	Headache
Angina	Sweating
Nervousness / Anxiety	Nausea / Vomiting
Tremors	Sore Throat

HOW SUPPLIED: Plastic Ampule of 2.5 mg / 3 mL (0.083% solution)
(TYPICALLY)

DOSAGES:

ADULT (>30 kg or 66 lbs) 2 ampules (5.0 mg) (may repeat once if needed)	PEDIATRIC (<30 kg or 66 lbs) 1 ampule (2.5 mg) (may repeat once if needed)
---	--

PEDIATRIC (<30 kg or 66 lbs)
1 mg IN (0.5 mg into each nostril) via Mucosal Atomization Device
(may repeat once if necessary to maintain respiratory activity)

ADULT (>30 kg or 66 lbs)
2 mg IN (1 mg into each nostril) via Mucosal Atomization Device
(may repeat once if necessary to maintain respiratory activity)

DOSAGES:

HOW SUPPLIED: Prefilled Syringe 2 mg / 2 mL
(TYPICALLY)

Tachycardia / Palpitations
Hypertension
Diaphoresis
Nausea / Vomiting

ADVERSE EFFECTS:

CAUTIONS: Naloxone may induce opiate withdrawal in patients who are physically opiate dependent

CONTRAINDICATIONS: None in the presence of respiratory depression or respiratory arrest

INDICATIONS: Unconscious with ineffective breathing and concerns for opiate use

Strong Adult Medical Control: (585) 271-2719
Strong Pediatric Medical Control: (585) 756-3430

NALOXONE (Narcan)

NITROGLYCERIN (Nitro)

Strong Adult Medical Control: (585) 271-2719

INDICATIONS: ACUTE CORONARY SYNDROME - Chest pain, suspicious of cardiac origin, including signs and symptoms of: retrosternal pain, heaviness, tightness or discomfort in the chest, radiation of pain or discomfort into jaw, arms or shoulders

CONTRAINDICATIONS: Systolic BP below 120 mmHg
Heart rate < 50 bpm or > 130 bpm
Medication not prescribed for the patient
In any patient who took medication for erectile dysfunction (Viagra™, Levitra™, Cialis™ or similar) within past 72 hours.

CAUTIONS: If systolic BP drops more than 20 mmHg, obtain medical consultation before further administration.

ADVERSE EFFECTS: Hypotension
Headache
Dizziness
Tachycardia

HOW SUPPLIED: 0.4 mg tablet
(TYPICALLY)

DOSAGES:	ADULT (>30 kg or 66 lbs) 1 tablet sublingual (under the tongue) (may repeat in 3-5 minutes if chest pain persists) MAX of 3 tablets (EMS doses + patient doses)	PEDIATRIC (<30 kg or 66 lbs) NOT INDICATED
-----------------	--	---

INDICATIONS: Signs and symptoms of an altered mental status resulting from hypoglycemia. For agencies approved to check finger stick blood glucose levels, a FSBG < 60 mm/dL is hypoglycemia (Any patient with a history of diabetes, presenting with an altered mental status, may be treated regardless of FSBG values.)

CONTRAINDICATIONS: Patient is unable to speak

CAUTIONS: All patients on oral hypoglycemic medications or long-acting insulin, who have been treated for potential hypoglycemia, should be transported. Treatment should not be withheld from patients with a stroke-like presentation, as this is likely due to hypoglycemia.

ADVERSE EFFECTS: None clinically significant

HOW SUPPLIED: 10-15 grams of glucose in various concentrations based on individual manufacturers. (TYPICALLY)

DOSAGES:

ADULT (>30 kg or 66 lbs) Administer 10-15 grams of glucose between the gum and cheek to conscious patient. (may require multiple smaller administrations)	PEDIATRIC (<30 kg or 66 lbs) Administer 10-15 grams of glucose between the gum and cheek to conscious patient. (may require multiple smaller administrations)
---	---

Strong Adult Medical Control: (585) 271-2719
Strong Pediatric Medical Control: (585) 756-3430

ORAL GLUCOSE